

AD-A196 055

4

CCS Research Report 591

THE CHANCE-CONSTRAINED CRITICAL PATH  
FOR A LARGE CLASS OF DISTRIBUTIONS


by

A. Charnes  
L. Gong

**CENTER FOR  
CYBERNETIC  
STUDIES**

The University of Texas  
Austin, Texas 78712

DTIC  
ELECTE  
JUL 13 1988  
S D  
E



This document has been approved  
for public release and sale in  
distribution is unlimited, except where  
indicated otherwise.

88 7 13 062 -

4

CCS Research Report 591

THE CHANCE-CONSTRAINED CRITICAL PATH  
FOR A LARGE CLASS OF DISTRIBUTIONS

by

A. Charnes  
L. Gong

March, 1988

The research for this paper was partly supported by National Science Foundation Grants SES 8520806 and U.S. Army Contract DAKF-15-87-C-0110, with the Center for Cybernetic Studies at The University of Texas at Austin, Austin, Texas 78712-1177. Reproduction in whole or in part is permitted for any purposes of the U.S. Government.

CENTER FOR CYBERNETIC STUDIES

A. Charnes, Director

College of Business Administration, 5.202  
The University of Texas at Austin  
Austin, Texas 78712-1177  
(512) 471-1821

DTIC  
ELECTE  
JUL 13 1988  
S D E

This document has been approved  
for public release and sale in  
distribution is unlimited.

CCS Research Report 591

THE CHANCE-CONSTRAINED CRITICAL PATH  
FOR A LARGE CLASS OF DISTRIBUTIONS

by

A. Charnes  
Linguo Gong

ABSTRACT

M. Kress proved for a special class of Location-Scale probability distributions there always exists a probability level for which the Chance Constrained Critical Path (CCCP) remains unchanged for all probabilities greater than or equal to that value. His chance constrained problem has zero-order decision rules and individual chance constraints.

This paper extends his results to most of the common probability distributions.

KEY WORDS

Chance Constrained Programming, (kp)

Chance Constrained Critical Paths

Location-Scale Distributions

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



Moshe Kress [1] studied the Chance Constrained Critical Path (CCCP) problem with zero order decision rule and individual chance constraints. He proved that for a class of Location-Scale probability distributions there always exists a probability level for which the CCCP remains unchanged for all probability values greater than or equal to that level. The purpose of this paper is to extend the results of [1] to most of the common probability distributions.

The CCCP problem with zero order decision rule and equal minimal probability level can be formulated as

$$P(\beta) \quad \begin{array}{ll} \text{Min } u_m - u_0 \\ \text{s.t.} \quad \left\{ \begin{array}{l} P_r \left( \sum_{i=0}^m \epsilon_{ij} u_i \geq t_j \right) \geq \beta \\ j = 1, 2, \dots, n \end{array} \right. \end{array}$$

where  $u_i$ 's are the start times of the activities.  $t_j$ ,  $j = 1, 2, \dots, n$ , are the durations of activities  $j$ 's which are the random variables with marginal distribution function  $F_j$ ;  $\epsilon_{ij}$  is the entry in row  $i$ , column  $j$  of the node-arc incidence matrix of the activity network.

$$\text{Let } F_j^{-1}(\beta) = \inf \{ t : F_j(t) \geq \beta \}$$

Notice that since the distribution function is right continuous, "inf" can be changed into "min". Now we can write  $P(\beta)$  into an equivalent deterministic form.

$$\begin{aligned}
 & \text{Min } u_m - u_0 \\
 P_1(\beta) \quad & \text{s.t.: } \begin{cases} \sum_{i=0}^m \epsilon_{ij} u_i \geq F_j^{-1}(\beta) \\ j = 1, 2, \dots, n \end{cases}
 \end{aligned}$$

The dual problem to  $P_1(\beta)$  is

$$\begin{aligned}
 & \text{Max } \sum_{j=1}^n F_j^{-1}(\beta) x_j \\
 D_1(\beta) \quad & \text{s.t.: } \begin{cases} \sum_{j=1}^n \epsilon_{ij} x_j = a_i, \quad i = 0, 1, \dots, m \\ x_j \geq 0, \quad j = 1, 2, \dots, n \end{cases}
 \end{aligned}$$

where  $a_0 = -1$ ,  $a_m = 1$ ,  $a_i = 0$ , for  $i = 1, 2, \dots, m-1$ .

Since  $D_1(\beta)$  is a pure network problem, for any given  $\beta \in (0, 1)$ , a basic optimal solution  $x^*(\beta) = \{x_1^*(\beta), x_2^*(\beta), \dots, x_n^*(\beta)\}$  of  $D_1(\beta)$  has the property  $x_j^*(\beta)$  is either 0 or 1, and  $\{j : x_j^*(\beta) = 1\}$  forms a critical path for the network of  $D_1(\beta)$ .

Let  $P = \{p_k : k = 1, 2, \dots, K\}$  denote the set of all paths from the source to the sink in the network, and  $J_k = \{j : \text{arc } j \text{ is in path } P_k\}$ . Then the problem  $P(\beta)$  is equivalent to the following problem.

$$\begin{aligned}
 & \text{Max} \sum_{k=1}^K \left( \sum_{j \in J_k} F_j^{-1}(\beta) \right) y_k \\
 \bar{P}(\beta) \quad & \text{s.t.} \quad \begin{cases} \sum_{k=1}^K y_k = 1 \\ y_k \geq 0 \\ k = 1, 2, \dots, K \end{cases}
 \end{aligned}$$

Clearly, if  $k^*$  such that

$$\sum_{j \in J_{k^*}} F_j^{-1}(\beta) = \text{Max}_{k \in I} \sum_{j \in J_k} F_j^{-1}(\beta)$$

where  $I = \{1, 2, \dots, K\}$ ,

then  $y_{k^*} = 1$ ,  $y_i = 0$  for  $i \neq k^*$  is an optimal solution of  $\bar{P}(\beta)$ , and if we set

$$x_j(\beta) = \begin{cases} 1 & \text{if } j \in J_{k^*} \\ 0 & \text{otherwise} \end{cases}$$

$x^*(\beta) = \{x_1^*(\beta), x_2^*(\beta), x_3^*(\beta), \dots, x_n^*(\beta)\}$  is an optimal solution of problem  $D_1(\beta)$ .

$$\text{Let } V_k(\beta) = \sum_{j \in J_k} F_j^{-1}(\beta),$$

$$U_{k,l}(\beta) = \sum_{j \in J_k \setminus J_l} F_j^{-1}(\beta),$$

$$H(\beta) = \left\{ k : V_k(\beta) = \text{Max}_{i \in I} V_i(\beta) \right\}.$$

The following theorem gives a sufficient condition under which the CCCP remains unchanged for all probabilities greater than or equal to a level  $\beta_0$ .

For the distribution function  $F_j$ , denote the support set of  $F_j$  ( $\text{supp}(F_j)$ ) as the interval

$$[\underline{\gamma}_j, \bar{\gamma}_j]$$

where  $\underline{\gamma}_j = \sup \{ \gamma : F_j(\gamma) = 0 \}$

$$\bar{\gamma}_j = \inf \{ \gamma : F_j(\gamma) = 1 \}$$

**Theorem:** Assume marginal distribution functions  $F_j, j = 1, 2, \dots, n$ , are continuous and the density functions  $f_j > 0$ , a.e., in  $\text{supp}(F_j)$ .

Case 1:  $\bar{\gamma}_j < \infty$ , for  $j = 1, 2, \dots, n$ .

If  $k^* \in H(1)$ , and there exists a  $0 \leq \hat{\beta} < 1$  such that

$$U'_{k^*,k}(\beta) \leq U'_{k,k^*}(\beta)$$

for any  $\beta \in [\hat{\beta}, 1]$  and  $k \in H(1)$ ,

then there exists a  $\beta_0 < 1$  such that for any  $\beta \in [\beta_0, 1]$

$$V_{k^*}(\beta) \leq V_k(\beta) \quad \text{for } k \in I$$

Case 2:  $\bar{\gamma}_j = +\infty$ , for  $j = 1, 2, \dots, n$ .

(a) If there exists a  $0 \leq \hat{\beta} < 1$  such that  $k^* \in H(\hat{\beta})$

$$\text{and } U'_{k^*,k}(\beta) \leq U'_{k,k^*}(\beta)$$

for any  $\beta \in [\hat{\beta}, 1]$  and  $k \in I$ .

or,

(b) if  $k^*$  is such that

$$\lim_{\beta \rightarrow 1^-} \frac{U'_{k^*,k}(\beta)}{U'_{k,k^*}(\beta)} > 1$$

for  $k \in I$

Then, there exists a  $\beta_0 < 1$  such that for any  $\beta \in [\beta_0, 1]$

$$V_{k^*}(\beta) \geq V_k(\beta) \quad \text{for } k \in I$$

Proof: Case 1.

Since  $F_j$  is a continuous monotone distribution function,  $f_j = F_j'$  is a Lebesgue measurable function. Since  $f_j > 0$  a.e.,  $1/f_j = (F_j^{-1})'$  is a Lebesgue measurable function. By the condition of theorem defining Case 1,  $U'_{k,k^*}(\beta) \geq U'_{k^*,k}(\beta)$  for any  $\beta \in [\hat{\beta}, 1]$  and  $k \in H(1)$ . Therefore, for any  $\beta \in [\hat{\beta}, 1]$  and  $k \in H(1)$

$$0 \leq \int_{\beta}^1 (U'_{k,k^*}(t) - U'_{k^*,k}(t)) dt$$

$$\begin{aligned}
&= \int_{\beta}^1 \left( \dot{V}_k(t) - \dot{V}_k^*(t) \right) dt \\
&= V_k(1) - V_k^*(1) - V_k(\beta) + V_k^*(\beta)
\end{aligned}$$

By hypothesis  $V_k(1) = V_{k^*}(1)$ , hence

$$V_{k^*}(\beta) \geq V_k(\beta) \quad \text{for any } k \in H(1)$$

If  $k \notin H(1)$ , then  $V_{k^*}(1) > V_k(1)$

and since  $f_j > 0$  a.e. in  $\text{supp}(F_j)$ ,  $F_j^{-1}$  is a strictly increasing continuous function and so is the sum of these which is  $V_k(\beta)$ . Therefore there exists a  $\beta_k < 1$  such that for any  $\beta \in [\beta_k, 1]$

$$V_{k^*}(\beta) \geq V_k(\beta).$$

Let  $\beta_0 = \max \{ \hat{\beta}, \beta_k \text{ for } k \in I \setminus H(1) \} < 1$

Then for  $\beta \in [\beta_0, 1]$ ,  $k \in I$

$$V_{k^*}(\beta) \geq V_k(\beta)$$

Case 2.a.

For any  $\beta \in [\hat{\beta}, 1]$  and  $k \in I$

$$\begin{aligned}
0 &\leq \int_{\beta}^{\hat{\beta}} \left( \dot{U}_{k^*,k}(t) - \dot{U}_{k,k^*}(t) \right) dt \\
&= \int_{\beta}^{\hat{\beta}} \left( \dot{V}_{k^*}(t) - \dot{V}_k(t) \right) dt \\
&= V_{k^*}(\hat{\beta}) - V_k(\hat{\beta}) - V_{k^*}(\beta) + V_k(\beta)
\end{aligned}$$

By hypothesis  $k^* \in H(\hat{\beta})$ , so we have

$$V_{k^*}(\beta) \geq V_k(\beta)$$

Therefore we can choose  $\beta_0$  at least equal to  $\hat{\beta}$ .

Case 2.b.

$$\text{Since } \lim_{\beta \rightarrow 1} \frac{U'_{k^*,k}(\beta)}{U'_{k,k^*}(\beta)} > 1$$



there exists a  $\hat{\beta}$  such that for any  $\beta \in [\hat{\beta}, 1]$

$$\frac{\dot{U}_{k^*,k}(\beta)}{\dot{U}_{k,k^*}(\beta)} \geq 1 + \epsilon$$

that is

$$\dot{U}_{k^*,k}(\beta) - \dot{U}_{k,k^*}(\beta) \geq \epsilon \dot{U}_{k,k^*}(\beta)$$

Hence

$$\begin{aligned} \epsilon \int_{\hat{\beta}}^{\beta} \dot{U}_{k,k^*}(t) dt &\leq \int_{\hat{\beta}}^{\beta} (\dot{U}_{k^*,k}(t) - \dot{U}_{k,k^*}(t)) dt = \int_{\hat{\beta}}^{\beta} (V_{k^*}(t) - V_k(t)) dt \\ &\leq V_{k^*}(\beta) - V_k(\beta) - (V_{k^*}(\hat{\beta}) - V_k(\hat{\beta})) \end{aligned}$$

Notice that if  $\bar{\gamma}_j = \infty$  and  $f_j > 0$  a.e. in  $\text{supp}(f_j)$  there exists a  $\beta_0 < 1$  such that

$$\int_{\hat{\beta}}^{\beta_0} \dot{U}_{k,k^*}(t) dt \geq (V_k(\hat{\beta}) - V_{k^*}(\hat{\beta})) / \epsilon$$

So, for any  $\beta > \beta_0$ ,

$$V_{k^*}(\beta) - V_k(\beta) \geq 0$$

Hence we have obtained a  $\beta_0$  which satisfies our requirement.

**Remark 1:** From the argument of our proof, it is clear that the theorem is also true when the  $F_j$ 's satisfy the assumed condition only in a interval of  $[\alpha, \bar{\gamma}]$ , where  $\alpha$  is a certain real value less than  $\bar{\gamma}$ .

**Remark 2:** When the set of random variables  $t_i$ ,  $i = 1, 2, \dots, n$  contains both finite support and infinite support distribution functions, then we can treat it as we did in the infinite support case.

We are now going to show that the location-scale distribution family which Kress [1] studied satisfies the hypotheses of our theorem.

The  $F_j$ ,  $j = 1, 2, \dots, n$ , for the Kress family are location-scale distributions with the same generating distribution  $\emptyset$ , that is,

$$F_j(x) = \emptyset\left(\frac{x - \mu_j}{\sigma_j}\right) \text{ for } j = 1, 2, \dots, n, \text{ where } \mu_j, \sigma_j \text{ are the location and scale parameters of } F_j$$

respectively.

The inverse of a location-scale distribution of a non-negative random variable is

$$F_j^{-1}(\beta) = \sigma_j \emptyset^{-1}(\beta) + \mu_j$$

where  $\emptyset^{-1}(\beta) \geq 0$  is the  $\beta$ th fractile of the generating distribution  $\emptyset$ .

$$\text{Let } S_k = \sum_{j \in J_k} \sigma_j \geq 0, \quad M_k = \sum_{j \in J_k} \mu_j \geq 0.$$

$$\text{Then } V_k(\beta) = S_k \emptyset^{-1}(\beta) + M_k$$

and the following corollary is obtained.

Corollary: Suppose  $F_j$ ,  $j = 1, 2, \dots, n$ , are such location-scale distributions,  $\emptyset$  a non-negative continuous distribution.

Case 1. Let  $\bar{\gamma} = \inf\{\gamma: \emptyset(\gamma) = 1\} < \infty$

If  $k^* \in I$  such that  $k^* \in H(1)$  and

$$M_{k^*} = \max_{j \in H(1)} M_j$$

Then, for any  $\beta \in [0, 1]$ ,  $k \in H(1)$

$$U_{k^*,k}(\beta) \geq U_{k,k^*}(\beta)$$

Case 2. Let  $\bar{\gamma} = +\infty$

If  $k^* \in I$  such that

$$S_{k^*} \geq S_k, \text{ for all } k \in I \text{ and } M_{k^*} \geq M_k, \text{ for all } k \in H, \text{ where } H = \left\{ i: S_i = \max_{j \in I} S_j \right\},$$

then there exists a  $\hat{\beta}$  such that for  $\beta \in [\hat{\beta}, 1]$ ,  $k \in H(\hat{\beta})$  and  $U_{k^*,k}(\beta) \geq U_{k,k^*}(\beta)$  for  $k \in I$ .

Proof: Case 1. Let  $k \in H(1)$ , then since

$$M_{k^*} \geq M_k \text{ and } V_{k^*}(1) = S_{k^*} \emptyset^{-1}(1) + M_{k^*} = S_k \emptyset^{-1}(1) + M_k = V_k(1),$$

we have  $S_{k^*} \leq S_k$ .

Therefore  $V_{k^*}(\beta) = S_{k^*}(\varnothing^{-1}(\beta)) \leq S_k(\varnothing^{-1}(\beta)) = V_k(\beta)$

That is  $U_{k^*,k}(\beta) \leq U_{k,k^*}(\beta)$  for  $k \in H(1)$

Case 2: By hypothesis, we know if  $k \in H$

$$V_{k^*}(\beta) = S_{k^*}\varnothing^{-1}(\beta) + M_{k^*} \geq S_k\varnothing^{-1}(\beta) + M_k = V_k(\beta)$$

that is,  $U_{k^*,k}(\beta) \geq U_{k,k^*}(\beta)$  (\*)

If  $k \in \bar{H}$ , then  $S_{k^*} > S_k$ . Let  $\mu = S_{k^*} - S_k > 0$ .

Since  $\bar{\gamma} = \infty$ , there is a  $\hat{\beta}$  such that for any  $\beta \in [\hat{\beta}, 1]$ ,  $\varnothing^{-1}(\beta) \geq \frac{M_k - M_{k^*}}{\mu}$

$$\begin{aligned} \text{Hence, } V_{k^*}(\beta) &= S_{k^*}\varnothing^{-1}(\beta) + M_{k^*} \\ &= S_k\varnothing^{-1}(\beta) + \mu\varnothing^{-1}(\beta) + M_{k^*} \\ &\geq S_k\varnothing^{-1}(\beta) + M_k = V_k(\beta) \end{aligned} \quad (**)$$

Combining (\*) and (\*\*), for  $\beta \in [\hat{\beta}, 1]$

$$U_{k^*,k}(\beta) \geq U_{k,k^*}(\beta)$$

On the other hand, because  $S_{k^*} \geq S_k$  for  $k \in I$ , we can directly conclude

$$V_{k^*}(\beta) = S_{k^*}(\varnothing^{-1}(\beta)) \geq S_k(\varnothing^{-1}(\beta)) = V_k(\beta)$$

which is equivalent to

$$U_{k^*,k}(\beta) \geq U_{k,k^*}(\beta)$$

Q.E.D.

Example 1:

Consider the project network depicted in Figure 1.




Figure 1

Let  $t_1 = t_2$  have distribution  $U(0, 1)$

$$\begin{array}{ll}
 t_3 \sim \text{density function} & f_3(x) = \begin{cases} \frac{1}{2}(2-x) & 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases} \\
 t_4 = t_5 \sim \text{density function} & f_4(x) = f_5(x) = \begin{cases} 2x & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases} \\
 t_6 = t_7 \sim \text{density function} & f_6(x) = f_7(x) = \begin{cases} 3x^2 & 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}
 \end{array}$$

Then  $\bar{\gamma}_i \leq 2 < \infty$

By solving the pure network problem  $D_1(1)$ , we get the optimal critical paths  $J_1 = \{0, 2, 1, 4\}$ ,

$J_2 = \{0, 2, 3, 4\}$ . See Figure 2.




Figure 2

Since  $U'_{1,2}(1) = \frac{1}{1} + \frac{1}{2} = \frac{3}{2}$   
 and  $U'_{2,1}(1) = \frac{1}{3} + \frac{1}{3} = \frac{2}{3}$

the critical path  $J_2$  is what we are seeking. By use of the optimal solution corresponding to  $J_2$  in  $D_1(\beta)$  and computing its reduced cost, we can get the minimal probability level  $\beta_0 = 0$ , for which CCCP remains unchanged for all probability values greater than or equal to it.

**Example 2:**

Consider the project network depicted in Figure 1 but let  $t_i \sim (1 - \alpha_i) F(x) + \alpha_i G(x)$  where  $F(x)$ ,  $G(x)$  are the distribution functions of  $N(0,1)$  and  $\text{Exp}(1)$  respectively and  $\alpha_i = \frac{1}{i+1}$ ,  $i = 1, 2, \dots, 7$ .

Since the possible critical paths are  $J_1 = \{0, 2, 1, 4\}$ ,  $J_2 = \{0, 2, 3, 4\}$ ,  $J_3 = \{0, 2, 4\}$ ,  $J_4 = \{0, 1, 4\}$ , we have

$$\lim_{\beta \rightarrow 1^-} \frac{U_{2,k}(\beta)}{U_{k,2}(\beta)} > 1 \quad \text{for } k = 1, 3, 5$$

So  $J_2$  is the desired critical path.

## REFERENCES

- [1] Kress, Moshe (1984) "The Chance Constrained Critical Path with Location-Scale Distribution," European Journal of Operational Research 18, pp. 359-363.
- [2] Charnes, A. and W.W. Cooper (1963), "Deterministic Equivalents for Optimizing and Satisficing Under Chance Constraints," Operations Research 11, pp. 18-39.
- [3] \_\_\_\_\_, \_\_\_\_\_ and G.L. Thompson, (1964), "Critical Path Analysis via Chance Constrained and Stochastic Programming," Operations Research 12, pp. 460-470.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER CCS 591	2. GOVT ACCESSION NO. A176 55	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  The Chance-Constrained Critical Path for a Large Class of Distributions		5. TYPE OF REPORT & PERIOD COVERED  Technical
		6. PERFORMING ORG. REPORT NUMBER CCS 591
7. AUTHOR(s)  A. Charnes, L. Gong		8. CONTRACT OR GRANT NUMBER(s)  Army DAKF15-87-C-0110 NSF SES 8520806
9. PERFORMING ORGANIZATION NAME AND ADDRESS Center for Cybernetic Studies Austin, Texas 78712		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS  USAREC Fort Sheridan, IL 60037		12. REPORT DATE April 1988
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  This document has been approved for public release and sale; distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Chance Constrained Programming, Chance-Constrained Critical Paths, Location-Scale Distributions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  M. Kress proved for a special class of Location-Scale probability distributions there always exists a probability level for which the Chance-Constrained Critical Path (CCCP) remains unchanged for all probabilities greater than or equal to that value. His chance constrained problem has zero-order decision rules and individual chance constraints. This paper extends his results to most of the common probability distributions.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE  
S/N 0102-014-6601

Unclassified  
SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)